

# An Accurate Quadrature for the Numerical Integration of Polynomial Functions

**Tahar Latrache**

Department of Civil Engineering, University of Tebessa, Tebessa, Algeria

**Email address:**

latrache.tahar@yahoo.ca

**To cite this article:**

Tahar Latrache. An Accurate Quadrature for the Numerical Integration of Polynomial Functions. *International Journal of Applied Mathematics and Theoretical Physics*. Vol. 4, No. 1, 2018, pp. 1-7. doi: 10.11648/j.ijamtp.20180401.11

**Received:** December 15, 2017; **Accepted:** January 8, 2018; **Published:** January 19, 2018

---

**Abstract:** The numerical integration of polynomial functions is one of the most interesting processes for numerical calculus and analyses, and represents thus, a compulsory step especially in finite elements analyses. Via the Gauss quadrature, the users concluded a great inconvenience that is processing at certain points which not required the based in finite element method points for deducting the form polynomials constants. In this paper, the same accuracy and efficiency as the Gauss quadrature extends for the numerical integration of the polynomial functions, but as such at the same points and nodes have chosen for the determination of the form polynomials. Not just to profit from the values of the polynomials at those points and nodes, but also from their first derivatives, the chosen points positions are arbitrary and the resulted deducted formulas are therefore different, as will be presented below and implemented.

**Keywords:** Integration Quadrature, Numerical Methods, Numerical Integration, Polynomial Functions, Accurate Methods

---

## 1. Introduction

The Gauss quadrature represents an accurate method for the numerical integration of polynomials and by a technique of subdividing the limit integral to rectangles and in some points called Gauss points, which represent in the parametric formulation, the coordinates of the ordinates of those rectangles. The classical method represents actually an exact numerical one for the integration of polynomials and very effective for any functions. In the literature, the Gauss quadrature is known as a very precise and effective numerical integration method [1, 21], and in finite elements some elementary matrices and vectors are defined as the numerically or the theoretically of some others; in structural mechanics, the elementary stiffness matrices are deducted via the Gauss quadrature of the scalar product of the elasticity by the deformation-displacement matrices and their transpose. The deformation-displacement matrices are in general computing only once, and used to compute stiffness and strains and therefore stresses matrices and vectors, the fact which in general calls to extrapolation functions to compute those quantities at the level of nodes, the main points used to express the form polynomials and thereof, the analysts needs to know the values of the

computed results at the level of.

In this new proposal, the developed method is addressed to the applied mathematics users and to the finite elements ones especially. Herein, the polynomial functions we proceed to integrate using this new approach, are defined such that their constants are related to the ordinates of these polynomials and their first derivatives at the chosen points to find the integral expression concerns and thus, the liberty to choose the number and the integral points positions is actually complete and anyway, following this method the developed formulas represent exactly the integrated polynomials with which, are related to the ordinates of these functions and their first derivatives at the points have chosen.

The approach methodology begins by fixing the parametric coordinates at the chosen points, deducting the polynomial functions constants and their first derivatives at those points in this formulation, and such that the numerical integrals are conducted in a parametric interval also, therefore the integral formulas will be also related to just those indicated values. The numerical formulation in Cartesian coordinates is deducted actually by a simple linear transformation. For the multiple integrals, of two or three dimensions, the process is doing by a multiplication

of the numerical parts of the formulas which called 'weights' and also by the ordinates or their first derivatives of multi-dimensions polynomial functions at the integral points.

Moreover, the method will be experimented by some examples and the results will be discussed.

$$P(x) = A_0 + A_1x + A_2x^2 + \dots + A_ix^i + \dots + A_{2n-1}x^{2n-1} \quad (1)$$

And its first derivative

$$P'(x) = A_1 + 2A_2x + \dots + iA_ix^{i-1} + \dots + (2n-1)A_nx^{2n-2} \quad (2)$$

In addition, to transform the polynomial function  $P(x)$  defined in the interval  $[x_1, x_2]$  in Cartesian coordinates to  $P(\xi)$  in  $[-1, +1]$ , the parametric ones, it is sufficient to relate the two coordinates by the expression

$$x = \frac{x_2 - x_1}{2} \xi + \frac{x_2 + x_1}{2} \quad (3)$$

Such that,  $P(x) = P(\xi)$  (Figure 1) and the integral can be written as

$$\int_{x_1}^{x_2} P(x) dx = \frac{x_2 - x_1}{2} \int_{-1}^{+1} P(\xi) d\xi \quad (4)$$

Expression with which,  $dx = \frac{x_2 - x_1}{2} d\xi$  and,

$$P(\xi) = a_0 + a_1\xi + a_2\xi^2 + \dots + a_i\xi^i + \dots + a_{2n-1}\xi^{2n-1} \quad (5)$$

The polynomial constants  $a_i$  can be easily deduced in relations with  $A_i$ , and with which  $L = x_2 - x_1$ , as follows

$$\begin{cases} a_0 = A_0 + \frac{L}{2}A_1 + \left(\frac{L}{2}\right)^2 A_2 + \dots + \left(\frac{L}{2}\right)^i A_i + \dots + \left(\frac{L}{2}\right)^{2n-1} A_{2n-1} \\ a_1 = \frac{L}{2}A_1 + 2\left(\frac{L}{2}\right)^2 A_2 + \dots + i\left(\frac{L}{2}\right)^i A_i + \dots + (2n-1)\left(\frac{L}{2}\right)^{2n-1} A_{2n-1} \\ \dots \\ a_i = \sum_{j=i}^{2n-1} \left(\frac{L}{2}\right)^j A_j \quad i = 0 \\ a_i = \left(\frac{L}{2}\right)^i A_i + \sum_{j=i+1}^{2n-1} j \left(\frac{L}{2}\right)^j A_j \quad 0 < i \leq 2n \\ \dots \end{cases} \quad (6)$$

Otherwise, in order to relate the constants  $a_i$  with  $\Phi(\xi_k)$  and  $\Phi'(\xi_k)$ , it is needed to express  $2n$  equations and  $n$  integral points in the parametric formulation. For a polynomial function of the first degree, one point sufficient, and of the third degree, of two points and so on

$$\begin{cases} P(\xi_k) = \Phi(\xi_k) = \Phi_k \\ P'(\xi_k) = \Phi'(\xi_k) = \Phi'_k \quad i = 1, 2, \dots, n, \quad \xi_k = 0, -1, +1, \dots \end{cases} \quad (7)$$

Expressions, such that the fixed points in the Cartesian coordinates shall be equal,  $x_k = \frac{L}{2}(\xi_k + 1) + x_1$

And,

$$\begin{cases} P(x_k) = \Phi\left(\frac{L}{2}(\xi_k + 1) + x_1\right) = \Phi(\xi_k) = \Phi_k \\ P'_{,\xi}(x_k) = \Phi'\left(\frac{L}{2}(\xi_k + 1) + x_1\right) = \frac{L}{2}\Phi'(\xi_k) = \frac{L}{2}\Phi'_k \end{cases} \quad (8)$$

1. Polynomial functions of the first degree

## 2. Expressions of the Polynomial Functions Constants

Suppose that  $\Phi(x)$  is a polynomial function defined and the proceeding is to express its integral in the interval  $[x_1, x_2]$ . The polynomial of the  $2n-1$  degree  $\Phi(x)$  can be exactly or approximately expressed by a polynomial function of the form




Figure 1. The two points representation of  $P(x)$  in the parametric formulation.

Suppose that the polynomial  $\Phi(x)$  is formed by a polynomial function  $P(x)$ , such that

$$P(x) = A_0 + A_1x$$

In the parametric coordinates, this polynomial function will be transform to the expression

$$\begin{cases} P(\xi) = \left(A_0 + \frac{L}{2}A_1\right) + \frac{L}{2}A_1\xi = a_0 + a_1\xi \\ P'_{,\xi}(\xi) = \frac{L}{2}A_1 = a_1 \end{cases} \quad \begin{cases} a_0 = A_0 + \frac{L}{2}A_1 + \left(\frac{L}{2}\right)^2 A_2 + \left(\frac{L}{2}\right)^3 A_3 \\ a_1 = \frac{L}{2}A_1 + 2\left(\frac{L}{2}\right)^2 A_2 + 3\left(\frac{L}{2}\right)^3 A_3 \\ a_2 = \left(\frac{L}{2}\right)^2 A_2 + 3\left(\frac{L}{2}\right)^3 A_3 \\ a_3 = \left(\frac{L}{2}\right)^3 A_3 \end{cases} \quad (13)$$

Thus, just only one integral point at the middle of the interval  $[-1 + 1]$  is necessary and sufficient to integrate exactly or approximately the function  $\Phi(\xi)$ , such that  $\xi_0 = 0$  and

$$\begin{cases} a_0 + a_1 \cdot 0 = \Phi_0 \\ a_1 = \Phi'_0 \end{cases}$$

Or,

$$\begin{cases} a_0 = \Phi_0 \\ a_1 = \Phi'_0 \end{cases} \quad (9)$$

Expressions, such that  $\Phi'$  is the derivative of  $\Phi$  with respect to  $\xi$ , we can therefore rewrite

$$P(\xi) = \Phi_0 + \Phi'_0 \xi \quad (10)$$

2. Polynomial functions of the third degree

The third degree polynomial functions can be given by the expression

$$P(x) = A_0 + A_1x + A_2x^2 + A_3x^3 \quad (11)$$

In Cartesian coordinates, and in parametric ones, by

$$P(\xi) = a_0 + a_1\xi + a_2\xi^2 + a_3\xi^3 \quad (12)$$

Such that,

$$P(\xi) = \frac{2(\Phi_1 + \Phi_{-1}) - (\Phi'_1 - \Phi'_{-1})}{4} + \frac{3(\Phi_1 - \Phi_{-1}) - (\Phi'_1 + \Phi'_{-1})}{4} \xi + \frac{(\Phi'_1 - \Phi'_{-1})}{4} \xi^2 + \frac{-(\Phi_1 - \Phi_{-1}) + (\Phi'_1 + \Phi'_{-1})}{4} \xi^3 \quad (16)$$

Following the same way, via the resolution of the equations system (7), the constants  $a_i$  of the polynomial function  $P(\xi)$  (5) can be then deducted and related to  $\Phi_k$  and  $\Phi'_k$  for superior degrees

### 3. The Integral Expressions of the Polynomial Functions

1. The first degree polynomials (one integral point)

For the first degree polynomial functions (10), their integral expression is the result of

$$I = \int_{-1}^{+1} P(\xi) d\xi = \left[ \Phi_0 \xi + \frac{\Phi'_0}{2} \xi^2 \right]_{-1}^{+1}$$

Or

$$I = 2 \times \Phi_0 \quad (17)$$

And in Cartesian coordinates, as such

$$\frac{L}{2} \times 2 \times \Phi_0 = L \times \Phi_0 = (x_2 - x_1) \times \Phi \left( x = \frac{x_2 + x_1}{2} \right) \quad (18)$$

Such that  $\Phi_0$  is the value of  $\Phi(x)$  at  $x = \frac{x_2 + x_1}{2}$  (i.e. at  $x = \frac{x_2 - x_1}{2} \xi + \frac{x_2 + x_1}{2}$ , and  $\xi = 0$ ).

2. The third degree polynomials (two integral points)

Based for the determination of  $a_i$  on the values of  $\Phi(\xi)$  and  $\Phi'(\xi)$  at  $-1$ . and  $+1$ . (i.e. at  $x_1$  and  $x_2$ ), the third degree polynomials are defined by the expression (16), and their integral in the parametric formulation

Moreover, in order to deduct the expressions relating between  $a_i$ ,  $\Phi_k$  and  $\Phi'_k$ , it shall resolve the linear equations system

$$\begin{cases} a_0 + a_1 + a_2 + a_3 = \Phi_1 \\ a_0 - a_1 + a_2 - a_3 = \Phi_{-1} \\ a_1 + 2a_2 + 3a_3 = \Phi'_1 \\ a_1 - 2a_2 + 3a_3 = \Phi'_{-1} \end{cases} \quad (14)$$

Such that  $\Phi(\xi)$  and  $\Phi'(\xi)$  are evaluated at  $\xi = +1$ . and  $\xi = -1$ . respectively (the two limits of the interval), the results can be given thus by

$$\begin{cases} a_0 = \frac{2(\Phi_1 + \Phi_{-1}) - (\Phi'_1 - \Phi'_{-1})}{4} \\ a_1 = \frac{3(\Phi_1 - \Phi_{-1}) - (\Phi'_1 + \Phi'_{-1})}{4} \\ a_2 = \frac{(\Phi'_1 - \Phi'_{-1})}{4} \\ a_3 = \frac{-(\Phi_1 - \Phi_{-1}) + (\Phi'_1 + \Phi'_{-1})}{4} \end{cases} \quad (15)$$

Therefore, the polynomial functions (12) in relation with the expressions of  $a_i$  (15), can be rewritten

$$I = \int_{-1}^{+1} P(\xi) d\xi = 2a_0 + \frac{2}{3}a_2 = 2 \frac{2(\Phi_1 + \Phi_{-1}) - (\Phi'_1 - \Phi'_{-1})}{4} + \frac{2(\Phi'_1 - \Phi'_{-1})}{3 \cdot 4}$$

Thus,

$$I = 1 \times (\Phi_1 + \Phi_{-1}) - \frac{1}{3} \times (\Phi'_1 - \Phi'_{-1}) \quad (19)$$

And in the Cartesian coordinates,

$$I = \int_{x_1}^{x_2} P(x) dx = \frac{(x_2 - x_1)}{2} \times \left[ (\Phi(x_2) + \Phi(x_1)) - \frac{1}{3} \times \frac{(x_2 - x_1)}{2} \times (\Phi'(x_2) - \Phi'(x_1)) \right] \quad (20)$$

Such that  $\Phi'$  is the first derivative of the function  $\Phi$  with respect to  $\xi$  and,

$$\frac{d\Phi(x)}{d\xi} = \frac{dx}{d\xi} \frac{d\Phi(x)}{dx} = \frac{L}{2} \frac{d\Phi(x)}{dx} \quad (21)$$

And the multiplication factor  $(x_2 - x_1)/2$  appears before, represents the Jacobean of the transformation, and appears also before the derivatives with respect to  $\xi$  according to (21).

3. The fifth degree polynomial functions (three integral points)

In this proposed approach, the liberty is considered complete for choosing the integral points positions, and with the Gaussian fixed points positions, the derivatives parts disappeared. Also, if equal distances between points is chosen, for instance the two interval limits and its middle, the constants  $a_0$ ,  $a_2$  and  $a_4$  are thus given by

$$\begin{aligned} a_0 &= \Phi_0 \\ a_2 &= \frac{4(\Phi_1 + \Phi_{-1}) - 8\Phi_0 - (\Phi'_1 - \Phi'_{-1})}{4} \\ a_4 &= \frac{-2(\Phi_1 + \Phi_{-1}) + 4\Phi_0 + (\Phi'_1 - \Phi'_{-1})}{4} \end{aligned}$$

$$I = W_1 \times (\Phi_1 + \Phi_{-1}) + W_2 \times (\Phi_{1/6} + \Phi_{-1/6}) + \overline{W}_1 \times (\Phi'_1 - \Phi'_{-1}) + \overline{W}_2 \times (\Phi'_{1/6} - \Phi'_{-1/6}) \quad (25)$$

Replacing  $\Phi_i$  and  $\Phi'_i$  by their expressions

$$I = W_1(2a_0 + 2a_2 + 2a_4 + 2a_6) + W_2 \left( 2a_0 + \frac{2}{6^2}a_2 + \frac{2}{6^4}a_4 + \frac{2}{6^6}a_6 \right) + \overline{W}_1(4a_2 + 28 + 16a_6) + \overline{W}_2 \left( \frac{14}{6^1}a_2 + \frac{2}{6^3}a_4 + \frac{2}{6^5}a_6 \right)$$

The two expressions (25) and (24) shall be equal, and such that the following equations system is resulted

$$\begin{cases} W_1 + W_2 = 1 \\ W_1 + \frac{1}{6^2}W_2 + 2\overline{W}_1 + \frac{2}{6}\overline{W}_2 = \frac{1}{3} \\ W_1 + \frac{1}{6^4}W_2 + 4\overline{W}_1 + \frac{4}{6^3}\overline{W}_2 = \frac{1}{5} \\ W_1 + \frac{1}{6^6}W_2 + 6\overline{W}_1 + \frac{6}{6^5}\overline{W}_2 = \frac{1}{7} \end{cases} \quad (26)$$

Therefore, with the same process and for the ninth degree polynomial functions (5 integral points), the described system is deduced

Thus, the integral of the fifth degree polynomials, and which related to  $a_0$ ,  $a_2$  and  $a_4$  by,

$$I = \int_{-1}^{+1} P(\xi) d\xi = 2a_0 + \frac{2}{3}a_2 + \frac{2}{5}a_4$$

Can be easily simplified by the expression,

$$I = \frac{7}{15} \times (\Phi_1 + \Phi_{-1}) + \frac{16}{15} \times \Phi_0 - \frac{1}{15} \times (\Phi'_1 - \Phi'_{-1}) \quad (22)$$

In parametric coordinates, and in the Cartesian ones by,

$$I = \int_{x_1}^{x_2} P(x) dx = \frac{(x_2 - x_1)}{2} \times \left[ \frac{7}{15} \times (\Phi(x_2) + \Phi(x_1)) + \frac{16}{15} \times \frac{x_2 + x_1}{2} - \frac{1}{15} \times \frac{(x_2 - x_1)}{2} \times (\Phi'(x_2) - \Phi'(x_1)) \right] \quad (23)$$

4. The seventh degree polynomial functions (four integral points)

Suppose that  $P(\xi)$  is a seventh degree polynomial function, its exact integral in the parametric coordinates is related to  $a_0$ ,  $a_2$ ,  $a_4$  and  $a_6$  by,

$$I = \int_{-1}^{+1} P(\xi) d\xi = 2a_0 + \frac{2}{3}a_2 + \frac{2}{5}a_4 + \frac{2}{7}a_6 \quad (24)$$

Otherwise, easily could be expressed as such,

$$\begin{cases} W_1 + 2W_2 + 2W_3 = 2 \\ W_2 + \frac{1}{4}W_3 + 2\overline{W}_1 + \overline{W}_2 = \frac{1}{3} \\ W_2 + \frac{1}{4^2}W_3 + 4\overline{W}_1 + \frac{1}{2}\overline{W}_2 = \frac{1}{5} \\ W_2 + \frac{1}{4^3}W_3 + 6\overline{W}_1 + \frac{3}{16}\overline{W}_2 = \frac{1}{7} \\ W_2 + \frac{1}{4^4}W_3 + 8\overline{W}_1 + \frac{3}{16}\overline{W}_2 = \frac{1}{9} \end{cases} \quad (27)$$

From one to five integral points, the weights  $W_i$  and  $\overline{W}_i$  of  $\Phi(\xi_i)$  and  $\Phi'(\xi_i)$  according to the number of points and their chosen positions  $\xi_i$  are cited in Table 1 (Appendix). Moreover, the lecturer can exercise polynomials of superior degrees or any favorable chosen points' positions.

Otherwise, in two dimensions it shall find the quadrature formulas for  $P = P(\xi, \eta)$  by integrating with respect to  $\xi$  and then with respect to  $\eta$ . For a quadrilateral interval area, the limit integral of the polynomial in the parametric formulation square area, would be formulated by

$$\begin{aligned} I &= \int_{-1}^{+1} \int_{-1}^{+1} P(\xi, \eta) d\xi d\eta = \int_{-1}^{+1} \left[ \sum_i W_i P(\xi_i, \eta) + \overline{W}_i P'_{,\xi}(\xi_i, \eta) \right] d\eta \\ &= \sum_j \left[ W_j \left[ \sum_i W_i P(\xi_i, \eta_j) + \overline{W}_i P'_{,\xi}(\xi_i, \eta_j) \right] + \overline{W}_j \left[ \sum_i W_i P'_{,\eta}(\xi_i, \eta_j) + \overline{W}_i P'_{,\xi\eta}(\xi_i, \eta_j) \right] \right] \end{aligned}$$

Or

$$I = \sum_i \sum_j [W_i W_j P_{ij} + \overline{W}_i W_j P'_{ij,\xi} + W_i \overline{W}_j P'_{ij,\eta} + \overline{W}_i \overline{W}_j P'_{ij,\xi\eta}] \quad (28)$$

Moreover, in three dimensions (or in a cubic area of a limit integral), the lecturer would be concluded then

$$I = \int_{-1}^{+1} \int_{-1}^{+1} \int_{-1}^{+1} P(\xi, \eta, \zeta) d\xi d\eta d\zeta$$

Is spreading by

$$I = \sum_i \sum_j \sum_k \left[ W_i W_j W_k P_{ijk} + \overline{W}_i W_j W_k P'_{ijk,\xi} + W_i \overline{W}_j W_k P'_{ijk,\eta} + W_i W_j \overline{W}_k P'_{ijk,\zeta} + \overline{W}_i \overline{W}_j W_k P'_{ij,\xi\eta} + W_i \overline{W}_j \overline{W}_k P'_{ijk,\eta\zeta} + \overline{W}_i \overline{W}_j \overline{W}_k P'_{ijk,\xi\zeta} + \overline{W}_i \overline{W}_j \overline{W}_k P'_{ij,\xi\eta\zeta} \right] \quad (29)$$

Expressions with which,  $\Phi_{ij}$  and  $\Phi'_{ij,\xi}$  are respectively, the ordinates of  $\Phi$  at the point  $(\xi_i, \eta_j)$  and its first derivative with respect to  $\xi$ , and so on for the remaining ordinates and their first derivatives. For more details, it is easier to determine the multidimensional integral formulas for polynomials defined in triangles and tetrahedrons.

### 4. Results and Discussion

For the element beam, with the elasticity modulus  $E$ , inertia moment  $I$  and length  $L$ , the deformation-displacement vector  $[B]$  is given by

$$[B] = \left[ -\frac{6}{L^2} + \frac{12}{L^3}x \quad -\frac{4}{L} + \frac{6}{L^2}x \quad \frac{6}{L^2} - \frac{12}{L^3}x \quad -\frac{2}{L} + \frac{6}{L^2}x \right]$$

And the stiffness matrix  $[k]$  shall be computed via integration

$$[k] = \int_0^L [B]^T EI [B] dx = EI \int_0^L [B]^T [B] dx$$

Using two integral points, for computing the stiffness element  $k_{11}$ , thus

$$k_{11} = EI \frac{L}{2} \left[ 1 \times (\Phi_1 + \Phi_{-1}) - \frac{1}{3} \times \frac{L}{2} (\Phi'_1 - \Phi'_{-1}) \right]$$

Expression such that,

$$\Phi(x) = \left( -\frac{6}{L^2} + \frac{12}{L^3}x \right)^2, \text{ and } \Phi'(x) = \frac{24}{L^3} \left( -\frac{6}{L^2} + \frac{12}{L^3}x \right)$$

Thus,

$$\Phi_1 = \frac{36}{L^4}, \Phi_{-1} = \frac{36}{L^4}, \Phi'_1 = \frac{144}{L^5} \text{ and } \Phi'_{-1} = -\frac{144}{L^5}$$

Via the expressed relation above, therefore

$$k_{11} = 12 \frac{EI}{L^3} = \text{exact solution}$$

Moreover, with the same way, the lecturer can easily verify the remaining elements of the matrix, and conclude that are just the constitution elements of the stiffness matrix for the two nodes beam element deduced by direct integration

$$[k] = \begin{bmatrix} 12 & 6L & -12 & 6L \\ & 4L^2 & -6L & 2L^2 \\ & & 12 & -6L \\ Sym & & & 4L^2 \end{bmatrix}$$

Exact results would be obtained for all the stiffness

elements, and using also the Gauss quadrature, putting  $x = \frac{L}{2}(\xi + 1)$  and  $\xi = \pm\sqrt{3}$ , and the between difference that the vector  $[B]$  is actually computed at  $\xi = \pm 1$  instead of  $\xi = \pm\sqrt{3}$ , the fact which conducting to compute the stresses and strains at the same points which represents an inconvenience with Gauss quadrature and in this contributed quadrature is actually modified.

For the cubic polynomial function

$$\Phi(x) = A_0 + A_1x + A_2x^2 + A_3x^3$$

Its direct exact integral from 0 to  $L$

$$\int_0^L \Phi(x) dx = A_0L + \frac{1}{2}A_1L^2 + \frac{1}{3}A_2L^3 + \frac{1}{4}A_3L^4$$

Using the numerical formula (20),

$$I = \frac{L}{2} \left[ (\Phi(L) + \Phi(0)) - \frac{1}{3} \frac{L}{2} (\Phi'(L) - \Phi'(0)) \right]$$

Thus,

$$\Phi(L) + \Phi(0) = 2A_0 + A_1L + A_2L^2 + A_3L^3$$

$$\Phi'(L) - \Phi'(0) = 2A_2L + 3A_3L^2$$

Replacing by in the formula above, we deduct

$$I = A_0L + \frac{1}{2}A_1L^2 + \frac{1}{3}A_2L^3 + \frac{1}{4}A_3L^4$$

Which is exactly the exact limit integral from 0 to  $L$  of  $\Phi(x)$ .

Using this approach, exact results would be obtained thus for polynomial functions with degrees less than the odd degree of  $\Phi(x)$  and approximately for every else mathematical functions, noting that the notion of the quadrature would be also applied to this proposition with respect to the ordinates of the polynomial functions themselves, with which the derivatives parts, only represent contributions to meet the integration exact results.

## 5. Conclusion

For the finite elements analyses, the stresses and strains and the elements of the deformation-displacement matrices in solid mechanics, the analysts in general are needing actually to obtain the results at the levels of nodes of elements, such that the extrapolation functions and especially at the edge nodes, don't allow to offer good precisions. Moreover, the proposed integration formulas for polynomials, which addressed especially to the finite elements developers, such as they could be more favorable for finite elements analysts, and also for the integration of polynomials in general. In addition, such as the same obtained numerical results using the developed formulas and would be otherwise, obtained using Gauss quadrature and exact direct integration ones, thereof the intervention of the first derivatives ordinates herein, mean just that the complete liberty of the choosing points positions, and their contributions in this effect to get

exact numerical integration of polynomial functions and avoid actually the required points' positions in the classical Gaussian quadrature.

## Appendix

**Table 1.** The scale of numbers, points' positions and the weights of the proposed quadrature.

$n$	$\xi_i$	$W_i(\Phi_i)$	$\bar{W}_i(\Phi'_i)$
1	0.	+2.	0.
2	-1.	+1.	+1/3
	+1.	+1.	-1/3
3	-1.	+7/15	+1/15
	0.	+16/15	0.
	+1.	+7/15	-1/15
4	-1.	+453457/1500625	+3419/128625
	-1/6	+1047168/1500625	-8352/42875
	+1/6	+1047168/1500625	+8352/42875
	+1.	+453457/1500625	-3419/128625
4	-1.	+61/189	+1/35
	-1/2	+128/189	+32/315
	+1/2	+128/189	-32/315
	+1.	+61/189	-1/35
5	-1.	+1601/8505	+29/2835
	-1/2	+4096/8505	-128/2835
	0.	+208/315	0.
	+1/2	+4096/8505	+128/2835
	+1.	+1601/8505	-29/2835

## References

- [1] Baldoni, V. N. Berline, De Loera, J. A. K'oppe, M. and Vergne M. (2010). How to integrate a polynomial over a simplex. *Mathematics of Computation*, DOI: 10.1090/S0025-5718-2010-02378-6.
- [2] Bernardini F. (1991) Integration of polynomials over n-dimensional polyhedra. *Comput Aided Des* 23 (1): 51–58.
- [3] Davis, P. J. and Rabinowitz, P. (1984a). *Methods of Numerical Integration*. Academic Press, San Diego, 2nd edition.
- [4] Davis, P. J. and Rabinowitz, P. (1984b). *Methods of Numerical Integration*. Academic Press, San Diego, 2nd edition.
- [5] Dunavant D. A. (1985) High degree efficient symmetrical Gaussian quadrature rules for the triangle. *Int J Numer Methods Eng* 21: 1129–1148.
- [6] Holdych D. J., Noble D. R., Secor R. B. (2008) Quadrature rules for triangular and tetrahedral elements with generalized functions. *Int J Numer Methods Eng* 73: 1310–1327.
- [7] Keast P. (1986) Moderate-degree tetrahedral quadrature formulas. *Comput Methods Appl Mech Eng* 55: 339–348.
- [8] Liu Y. and Vinokur M. (1998). Exact integrations of polynomials and symmetric quadrature formulas over arbitrary polyhedral grids. *Journal of Computational Physics*, 140: 122–147.
- [9] Lubinsky, D. S. and Rabinowitz, P. (1984). Rates of convergence of Gaussian quadrature for singular integrands. *Mathematics of Computation*, 43 (167): 219–242. Press.

- [10] Lyness J. N., Jespersen D. (1975) Moderate degree symmetric quadrature rules for the triangle. *J Inst Math Appl* 15: 19–32.
- [11] Lyness J. N., Monegato G. (1977) Quadrature rules for regions having regular hexagonal symmetry. *SIAM J Numer Anal* 14 (2): 283–295.
- [12] Mousavi S. E., Xiao H., Sukumar N. (2010) Generalized Gaussian quadrature rules on arbitrary polygons. *Int J Numer Methods Eng* 82 (1): 99–113.
- [13] Mousavi S. E., Sukumar N. (2010) Generalized Duffy transformation for integrating vertex singularities. *Comput Mech* 45 (2–3): 127–140.
- [14] Mousavi S. E., Sukumar N. (2010) Generalized Gaussian quadrature rules for discontinuities and crack singularities in the extended finite element method. *Comput Methods Appl Mech Eng* 199 (49–52): 3237–3249.
- [15] Rathod H. T., Govinda Rao H. S. (1997) Integration of polynomials over n-dimensional linear polyhedra. *Comput Struct* 65 (6): 829–847
- [16] Silvester P. (1970) Symmetric quadrature formulae for simplexes. *Math Comput* 24 (109): 95–100.
- [17] Sunder K. S., Cookson R. A. (1985) Integration points for triangles and tetrahedrons obtained from the Gaussian quadrature points for a line. *Comput Struct* 21 (5): 881–885.
- [18] Ventura G. (2006) On the elimination of quadrature subcells for discontinuous functions in the eXtended finite-element method. *Int J Numer Methods Eng* 66: 761–795.
- [19] Wandzura S., Xiao H. (2003) Symmetric quadrature rules on a triangle. *Comput Math Appl* 45: 1829–1840.
- [20] Wong, R. (1989). *Asymptotic approximation of integrals*. Academic Press, San Diego.
- [21] Xiao H., Gimbutas Z. (2010) A numerical algorithm for the construction of efficient quadratures in two and higher dimensions. *Comput Math Appl* 59: 663–676.